

Polyborate anions templated by cationic transition-metal complexes:

Altahan, Mohammed; Beckett, Michael; Coles, Simon J.; Horton, Peter N.

Phosphorus, Sulfur and Silicon and the Related Elements

DOI:

[10.1080/10426507.2015.1128907](https://doi.org/10.1080/10426507.2015.1128907)

Published: 01/04/2016

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Altahan, M., Beckett, M., Coles, S. J., & Horton, P. N. (2016). Polyborate anions templated by cationic transition-metal complexes: $[\text{Co}(\text{diNOsar})]_2[\text{B}_3\text{O}_3(\text{OH})_4]\text{Cl}_{5.4.75} \cdot \text{H}_2\text{O}$. *Phosphorus, Sulfur and Silicon and the Related Elements*, 191(4), 572-575.
<https://doi.org/10.1080/10426507.2015.1128907>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Polyborate anions templated by cationic transition-metal complexes: $[\text{Co}(\text{diNOsar})]_2[\text{B}_3\text{O}_3(\text{OH})_4]\text{Cl}_5 \cdot 4.75 \text{H}_2\text{O}$

Mohammed A. Altahan¹, Michael A. Beckett^{1*}, Simon J. Coles², Peter N. Horton²

¹School of Chemistry, Bangor University, Bangor, Gwynedd, LL57 2UW, UK

²School of Chemistry, Southampton University, Southampton, SO17 1BG, UK

GRAPHICAL ABSTRACT



Abstract The transition metal complexes $[\text{Co}(\text{diNOsar})]^{3+}$, $[\text{Co}(\text{en})_3]^{3+}$, $[\text{Co}(\text{dien})_2]^{3+}$ have been used to template unusual isolated polyborate anions by reaction of basic aqueous solutions containing the cations with $\text{B}(\text{OH})_3$ (1:10). The synthesis and characterization (TGA/DSC, IR, NMR, single-crystal XRD) of one such complex $[\text{Co}(\text{diNOsar})]_2[\text{B}_3\text{O}_3(\text{OH})_4]\text{Cl}_5 \cdot 4.75\text{H}_2\text{O}$ (**1**) is reported. Compound **1** is a rare example of a salt containing the isolated triborate(1-) anion, and its formation is engineered by the cation, which has extensive H-bond interactions with the anion in its secondary coordination sphere.

Keywords Triborate(1-); Self-assembly; H-bonds; $[\text{Co}(\text{diNOsar})]^{3+}$; X-ray Structure; Oxidoborate

Received xx yyyy 2015; accepted xx yyyy 2015

Acknowledgements: We thank the EPSRC for use of the NCS Service.

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423520. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

INTRODUCTION

Polyborate anions are conveniently classified¹ as either polymeric (2D or 3D chains, sheets or networks) or discrete ‘isolated’ anionic moieties. Isolated polyborate anions are readily synthesized by the addition of B(OH)₃ to a basic aqueous solution containing templating cations², or by solvothermic³ methods. Templated products arise since B(OH)₃ in basic aqueous solution exists as a dynamic combinatorial library of polyborate anions⁴. In general, pentaborate(1-) salts (containing the [B₅O₆(OH)₄]⁻ anion) are formed because these salts have a strong H-bonded anionic lattice, which is sufficiently flexible to accommodate many medium sized cations⁵. Salts containing polyborate anions other than pentaborate(1-) are relatively rare⁶. We are interested in the synthesis of structurally novel polyborate anions and have adopted a strategy of templating such species by the use of sterically demanding and/or highly charged cations^{7,8}. Herein, we report the synthesis of several salts containing isolated polyborate anions partnered with transition metal cations and describe in detail the synthesis and characterization of the salt [Co(diNOsar)]₂[B₃O₃(OH)₄]Cl₅·4.75 H₂O (**1**) which contains the isolated triborate(1-) anion.

RESULTS AND DISCUSSION

Synthesis

Halide salts of transition-metal complexes are readily transformed, provided they are stable in basic solution, into their hydroxide salts by use of a Dowex 550A hydroxide activated anion exchange resin. The resulting aqueous solutions, which are separated from the resin, were reacted with MeOH/H₂O solutions of B(OH)₃ (1:10) to afford polyborate anions salts of cationic transition-metal complexes. Several salts have been prepared *e.g.*

[Co(diNOsar)]₂[B₃O₃(OH)₄]Cl₅·4.75H₂O (**1**), [Co(en)₃][B₅O₆(OH)₄][B₈O₁₀(OH)₆]·5H₂O (**2**), *s-fac*-[Co(dien)₂][B₇O₉(OH)₆]·9H₂O (**3**). Compounds **2** and **3**, which contain previously unobserved octaborate(2-) and heptaborate(3-) anions, were prepared in reasonable yields (~40%) and details^{7,8} of their synthesis and full characterization can be found elsewhere. The starting complex [Co(diNOsar)]Cl₃ was prepared by a reported⁹ procedure for the synthesis of **1**. Compound **1** was isolated in low yield (10%) and its high chloride content would indicate that

the ion exchange was not completely successful in this case. Nevertheless, we were able to characterize¹⁰ the compound by elemental analysis, mpt, thermally (TGA/DSC) and by NMR (¹H, ¹³C, ¹¹B) and IR spectroscopy. Single-crystals were also suitable for a XRD study.

NMR spectra of **1** were obtained in D₂O. The ¹H NMR spectrum of **1** consists of AB doublets for the CH₂ groups of the capping units and a complex AA'BB' pattern¹¹ (looking like a doublet with *J* = 9.2 Hz) for the CH₂ protons of the ethylenediamine rings. The secondary amine protons are observed with the residual HOD signal at δ = 4.70 ppm. The ¹³C{¹H} spectrum shows the expected 3 signals at δ = 51.4 (ethylene diamine groups), δ = 55.1 (CH₂ of capping groups) and δ = 87.8 ppm (tertiary C-NO₂ cap). As a consequence of boron exchange associated with rapid B(OH)₃/[B(OH)₄]⁻/polyborate equilibria⁴ the ¹¹B spectrum shows just one signal at δ = +16.2 ppm.

Thermal (TGA) analysis in air was consistent with a three step process involving low temperature (<125 °C) loss of water of crystallization, further dehydration to form an anhydrous polyborate (200-275 °C), and oxidation of organics (275-800 °C) to leave a glassy inorganic residue of composition Co₂B₃O₅Cl₅. Such thermal decomposition has been previously observed¹² for transition-metal polyborate salts.

X-ray structure of [Co(diNOsar)]₂[B₃O₃(OH)₄]Cl₅·4.75H₂O

Full crystallographic data for **1** are available in the supplementary information¹³ and crystallographic data are summarized in a footnote¹⁴. Compound **1** is an ionic compound and is a rare example of a salt containing the triborate(1-) anion. In addition to the [B₃O₃(OH)₄]⁻ anion, **1** contains 2 independent [Co(diNOsar)]³⁺ cations, 5 Cl⁻ anions and 4.75 H₂O molecules. There is a disordered water within the asymmetric unit. One area can exist as either one (25%) or two (75%) molecules of water, hence the fractional formula. Atomic labelling for the triborate(1-) anion is shown in Figure 1.

The [Co(diNOsar)]³⁺ cation comprises a central Co³⁺ ion surrounded by the 6 N donor atoms of the hexadentate neutral 'diNOsar' ligand in a distorted octahedral environment. *Cis* and *trans* N-Co-N angles range from 86.43(9) - 92.58(9)° (av. 90.06°) and 175.79(9) - 178.12(10)°

Received xx yyyy 2015; accepted xx yyyy 2015

Acknowledgements: We thank the EPSRC for use of the NCS Service.

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423520. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

(av. 176.62°) respectively. The Co-N bond lengths range from 1.963(2) - 1.975(2) Å, av. 1.969 Å. These data are not significantly different from previously reported data¹⁵ for this cation.

The $[\text{B}_3\text{O}_3(\text{OH})_4]^-$ ion of **1** possesses a B_3O_3 (boroxole) ring with one 4-coordinate and two 3-coordinate B centres. B-O bond lengths to 3-coordinate (B2 and B3) and 4-coordinate (B1) B centres range from 1.347(4) - 1.386(4) Å (av. 1.363 Å) and 1.466(4) - 1.471(4) Å (av. 1.488 Å), respectively. B2-O1 (1.351(3) Å) and B3-O3 (1.347(4) Å) are significantly shorter than the other B-O bonds within the boroxole rings. The boroxole ring is planar with ring angles ranging from 112.3(2) – 122.4(2)°. The smallest ring angle (O1B1O3) involves the 4-coordinate B centre, whereas ring angles at O1, O2, and O3 average 121.1° and those at the 3-coordinate B centres (B2 and B3) average 121.6°. This pattern of bond angles and bond lengths are in accord with previously reported data for this anion¹⁶ and mirror corresponding data observed¹⁷ in related isoelectronic systems such as $\text{Ar}_3\text{B}_3\text{O}_3\text{L}$ and $[\text{Ph}_3\text{B}_3\text{O}_3(\text{OH})]^-$.




Figure 1 Cation-triborate H-bond interactions in **1**. $d(\text{N13-H13})$ 1.00 Å, $d(\text{H13}\cdots\text{O4})$ 1.98 Å, $d(\text{N13}\cdots\text{O4})$ 2.935(3) Å, $\angle(\text{N13H13O4})$ 159.2°; $d(\text{N16-H16})$ 1.00 Å, $d(\text{H16}\cdots\text{O4})$ 1.73 Å, $d(\text{N16}\cdots\text{O4})$ 2.731(3) Å, $\angle(\text{N16H16O4})$ 177.4°; $d(\text{N32H32})$ 1.00 Å, $d(\text{H32}\cdots\text{O5})$ 1.81 Å, $d(\text{N32}\cdots\text{O5})$ 2.810(3) Å, $\angle(\text{N32H32O5})$ 175.8°; $d(\text{N35-H35})$ 1.00 Å, $d(\text{H35}\cdots\text{O5})$ 1.93 Å, $d(\text{N35}\cdots\text{O5})$ 2.898(3) Å, $\angle(\text{N35H35O5})$ 163.4°.

Received xx yyyy 2015; accepted xx yyyy 2015

Acknowledgements: We thank the EPSRC for use of the NCS Service.

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423520. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

There are multiple H-bond interactions within the solid-state structure of **1**. Full details can be found in the supplementary information. Polyborate anions usually display multiple anion-anion H-bond interactions and these are believed⁵ to be the driving force in the synthesis of non-metal cation pentaborate(1-) salts. Anion-anion interactions were found to be present in **2** and **3**. However, in **1**, triborate-triborate H-bond interactions are absent and each triborate(1-) anion is isolated from others by H₂O bridges, forming a sheet-like structure (Figure 2). This isolation implies that the crystal structure of **1** is engineered by interactions with other components within the system. The triborate(1-) anion is able to form 4 donor H-bonds and has 7 potential H-bond acceptor sites. The triborate(1-) anion in **1** does indeed form 4 donor interactions: three to H₂O molecules and one to a Cl⁻ anion (O6H6...Cl2). The triborate(1-) anion accepts 2 H-bonds at each hydroxyl group (O4 and O5) on the 4-coordinate B1, and also at O6 from H₂O (O54-H54...O6). The H-bonds directed at O4 and O5 originate from amine hydrogens on two [Co(diNOsar)]³⁺ cations in a chelating mode, as illustrated in Figure 1. These H-bond interactions are likely to be responsible for templating this salt and assembling the triborate(1-) anion. Secondary amine hydrogens of [Co(diNOsar)]³⁺ are known¹⁵ to form similar H-bond interactions *e.g.* [Co(diNOsar)][S₂CN(CH₂)₄]₃.




Figure 2 The triborate(1-) anions are arranged in planes linked together by H-bonds from H₂O molecules. The [Co(diNOsar)]³⁺ cations fill the cavities and H-bond to the triborate(1-) anions.

Received xx yyyy 2015; accepted xx yyyy 2015

Acknowledgements: We thank the EPSRC for use of the NCS Service.

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423520. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Summary

[Co(diNOsar)]₂[B₃O₃(OH)₄]Cl₅·4.75H₂O is a rare example of a salt containing the isolated triborate(1-) anion. It has been obtained from a self-assembly reaction of (partially) OH⁻ exchanged aqueous solution of [Co(diNOsar)]Cl₃ with B(OH)₃ (1:10). The solid-state structure of **1** shows the anion in the secondary coordination sphere of the cation. There are strong cation-anion H-bond interactions, which help to engineer the formation of the triborate(1-) anion.

REFERENCES

1. (a) Christ, C.L.; Clarke, J.R. *Phys Chem. Miner.* **1977**, 2, 59-87; (b) Schubert, D.M.; Smith, R.A.; Visi, M.Z. *Glass Technol.* **2003**, 44, 63-70.
2. Beckett, M.A.; Horton, P.N.; Coles, S.J.; Martin, D.W. *Inorg. Chem.*, **2011**, 50, 12215-12218.
3. (a) Wang, G.-M.; Pan, C.-Y.; Zheng, S.-T.; Yang, G.-Y. *Acta Cryst.* **2007**, E63, o1104-o1105; (b) Wang, G.-M.; Sun Y.-Q.; Yang, G.-Y. *J. Solid State Chem.* **2004**, 177, 4648-4654.
4. (a) Salentine, C.G. *Inorg. Chem.* **1983**, 22, 3920-3924; (b) Anderson, J.L.; Eyring, E.M.; Whittaker, M.P. *J. Phys. Chem.* **1964**, 68, 1128-1132.
5. Beckett, M.A.; Coles, S.J.; Davies, R.A.; Horton, P.N.; Jones, C.L. *Dalton Trans.* **2015**, 44, 7032-7040.
6. Schubert, D.M.; Knobler, C.B. *Phys. Chem. Glasses: Eur. J. Glass Sci. Technol. B* **2009**, 50, 71-78.
7. Altahan, M.A.; Beckett, M.A.; Coles, S.J.; Horton, P.N. *Inorg. Chem.* **2015**, 54, 412-414.
8. Altahan, M.A.; Beckett, M.A.; Coles, S.J.; Horton, P.N. *Inorg. Chem. Commun.* **2015**, 59, 95-98.
9. Geue, R.J.; Hambley, T.W.; Harrowfield, J.M.; Sargeson, A.M.; Snow, M.R. *J. Am. Chem. Soc.* **1984**, 106, 5478-5488.
10. Characterization data for **1**: C₂₈H_{73.50}B₃Cl₅Co₂N₁₆O_{19.75}. Elem. Anal Calc: C, 26.3; H 5.8; N, 17.5. Found: C, 25.6; H 6.5; N, 16.7%. Mp 285 °C. TGA: - 4.75 H₂O 7.8 % (6.7 % calc.), - 2H₂O 3.2 % (2.8% calc.), cation oxidation 57.7 % (58.6% calc.), residual Co₂B₃O₃Cl₅ 31.3% (31.9 % calc). IR (KBr/cm⁻¹): 3401(br,s), 3041(vs), 2871(vs), 1558(vs), 1432(m), 1345(s), 1077(m), 988(w), 849(w), 813(w). NMR (400 MHz, D₂O) δ¹H/ppm: 3.77 6H (d, 13.8 Hz), 3.41 6H (m), 3.24 6H (d, 13.8 Hz), 2.83 6H (m). δ¹¹B/ppm: 16.2. δ¹³C/ppm: 87.8, 55.1, 51.4.
11. Bernhardt, P.V.; Bygott, A.M.T.; Geue, R.J.; Hendrey, A.J.; Korybut-Daszkiewicz, B.R.; Lay, P.A.; Pladziejewicz, J.R.; Sargeson, A.M.; Willis, A.C. *Inorg. Chem.* **1994**, 33, 4553-4561.
12. (a) Wang, G.-M.; Sun, Y.-Q.; Yang, G.-Y. *J. Solid State Chem.* **2006**, 179, 1545-1553; (b) Liu, Z.-H.; Zhang, J.-J.; Zhang, W.-J. *Inorg. Chim. Acta* **2006**, 359, 519-524; (c) Yang, Y.; Wang, Y.; Zhu, J.; Liu, R.-B.; Xu, J.;

Received xx yyyy 2015; accepted xx yyyy 2015

Acknowledgements: We thank the EPSRC for use of the NCS Service.

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423520. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

- Meng, C.-C. *Inorg. Chim. Acta* **2011**, 376, 401-407; (d) You, X.; Zhu, L.; Sun, J. *Chin. J. Chem.* **2010**, 28, 2174-2178.
13. (a) Coles, S.J.; Gale, P.A. *Chem. Sci.* **2012**, 3, 683-689; (b) Rigaku, CrystalClear-SM Expert 3.1 **2013**, b27; (c) Sheldrick, G.M. *Acta Crystallogr.* **2015**, C71, 3; (d) Farrugia, L.J.; *J. Appl. Crystallogr.* **1997**, 30, 564; (e) Palatinus, L.; Chapuis, G.; *J. Appl. Crystallogr.* **2007**, 40, 786-790; (f) CrysAlisPro, Agilent Technologies, version **2014**, 1.171.37.31.
14. A Rigaku AFC12 goniometer equipped with an enhanced sensitivity (HG) Saturn724+ detector mounted at the window of a FR-E+ SuperBright molybdenum rotating anode generator with VHF Varimax optics (70 μ m focus) was used. X-ray data for **1**: C₂₈H_{73.50}B₃Cl₅Co₂N₁₆O_{19.75}, M = 1278.06; triclinic space group *P*-1, *a* = 9.6900(4) Å, *b* = 10.0170(6) Å, *c* = 29.1374(7) Å, α = 85.079(3) $^\circ$, β = 88.439(3) $^\circ$, γ = 62.137(5) $^\circ$; *U* = 2490.9(2) Å³; *Z* = 2; *T* = 100K; Wavelength = 0.71075 Å; density (calc) = 1.704 Mg/m³; Absorption coefficient 1.024 mm⁻¹; Crystal: orange lath, 0.180 x 0.030 x 0.010 mm³; *F*(000) 1331; θ range for data collection 2.420-29.944 $^\circ$; Index ranges -13 $\leq h \leq$ 12, -13 $\leq k \leq$ 13, -38 $\leq l \leq$ 40; 36914 reflections collected with 12826 independent reflections [*R*_{int} = 0.0478]; Completeness to θ = 25.242 $^\circ$, 99.9%; absorption correction: semi-empirical from equivalents; refinement method: full-matrix least squares on *F*²; data/restraints/parameters 12826/0/668; goodness-of-fit on *F*², 1.023; Final *R* indices [*F*² > 2 σ (*F*²)] *RI* = 0.0485, *wR*2 = 0.0976; *R* indices (all data) *RI* = 0.0740, *wR*2 = 0.1072; extinction coefficient n/a; largest diff. peak and hole 0.881 and -0.662 e Å⁻³.
15. Achilleos, A.A.; Gahan, L.R.; Hambley, T.W.; Healy, P.C.; Weedon, D.M. *Inorg. Chim. Acta* **1989**, 157, 209-214.
16. (a) Negro, A.D.; Pozaz, J.M.M.; Ungaretti, L. *Am. Mineral* **1975**, 60, 897-883; (b) Salentine, G.C.; *Inorg. Chem.* **1987**, 26, 128-132; (c) Beckett, M.A.; Horton, P.N.; Hursthouse, M.B.; Timmis, J.L. *RSC Adv.* **2013**, 3, 15185-15191; (d) Schubert, D.M.; Visi, M.Z.; Knobler, C.B. *Inorg. Chem.* **2008**, 47, 2017-2023.
17. (a) Beckett, M.A.; Coles, S.J.; Light, M.E.; Fischer, L.; Stiefvater-Thomas, B.M.; Varma, K.S. *Polyhedron* **2006**, 25, 1011-1016; Beckett, M.A.; Hibbs, D.E.; Hursthouse, M.B.; Owen, P.; Malik, K.M.A.; Varma, K.S. *Main Group Chem.* **1998**, 2, 251-258; (c) Beckett, M.A.; Strickland, G.C.; Varma, K.S.; Hibbs, D.E.; Hursthouse, M.B.; Malik, K.M.A. *J. Organomet. Chem.* **1997**, 535, 33-41.

Received xx yyyy 2015; accepted xx yyyy 2015

Acknowledgements: We thank the EPSRC for use of the NCS Service.

Address correspondence to Dr M.A. Beckett, School of Chemistry, Bangor University, Bangor, Gwynedd, UK.

Email: m.a.beckett@bangor.ac.uk

Electronic supplementary information (ESI) available. CCDC 1423520. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

POLYBORATE ANIONS TEMPLATED BY METAL COMPLEXES